

Of Note: this is the post peer-reviewed version. Printed pdf is found on the publisher website of Human Vaccines & Immunotherapeutics.

Dendritic cell immunotherapy in uterine cancer

An Coosemans^{1,2,†,*}, Sandra Tuyaerts^{3,†}, Anke Vanderstraeten³, Ignace Vergote^{1,2}, Frédéric Amant^{1,2}, and Stefaan W Van Gool^{4,5}

¹Department of Oncology; Leuven Cancer Institute; KU Leuven; Leuven, Belgium

²Department of Gynecology and Obstetrics; UZ Leuven; Leuven, Belgium

³Department of Oncology; Division of Gynecologic Oncology; KU Leuven, Belgium

⁴Department of Microbiology and Immunology; KU Leuven, Leuven, Belgium

⁵Department of Pediatrics; UZ Leuven, Leuven, Belgium

[†]These authors contributed equally to this work.

Keywords: uterine cancer, dendritic cell, DC, immunotherapy, WT1, electroporation

Abbreviations: AICD, activation-induced cell death; CCR, C-C chemokine receptor; CCL, chemokine ligand; CD, cluster of differentiation; CD40L, cluster of differentiation 40 ligand; DC, dendritic cell; DC-LAMP, dendritic cell lysosomal-associated membrane protein; FIGO, international federation of gynecology and obstetrics; GM-CSF, granulocyte macrophage colony-stimulating factor; HLA, human leukocyte antigen; hTERT, human telomerase reverse transcriptase; IL, interleukin; LPS, lipopolysaccharide; MCP, monocyte chemoattractant protein 1; MDSC, myeloid derived suppressor cell; MHC, major histocompatibility complex; MIP, macrophage inflammatory protein; mRNA, messenger ribonucleic acid; MUC1, mucin-1; RNA, ribonucleic acid; TAA, tumor-associated antigen; TAM, tumor-associated macrophage; TGF β , tumor growth factor beta; TIL, tumor infiltrating lymphocyte; TLR, toll-like receptor; Treg, regulatory T cell; TNF α , tumor necrosis factor alpha; WT1, Wilms' tumor gene 1

*Correspondence to: An Coosemans; Email: an.coosemans@gmail.com

Submitted: 03/20/2014; Accepted: 03/31/2014

<http://dx.doi.org/10.4161/hv.28716>

Uterine cancer is the most common pelvic gynecological malignancy. Uterine sarcomas and relapsed uterine carcinomas have limited treatment options. The search for new therapies is urgent. Dendritic cell (DC) immunotherapy holds much promise, though has been poorly explored in uterine cancer. This commentary gives an insight in existing DC immunotherapy studies in uterine cancer and summarizes

the possibilities and the importance of the loading of tumor antigens onto DC and their subsequent maturation. However, the sole application of DC immunotherapy to target uterine cancer will be insufficient because of tumor-induced immunosuppression, which will hamper the establishment of an effective anti-tumor immune response. The authors give an overview on the limited existing immunosuppressive data and propose a novel approach on DC immunotherapy in uterine cancer.

Introduction

Worldwide, uterine cancer is the seventh most common malignant disorder and the most frequent pelvic gynecological cancer. Each year, in Europe, an estimated 9000 women die of the disease. It occurs mainly in postmenopausal women with a mean age of 65 y.

Uterine tumors can be epithelial or mesenchymal in origin and are designated as carcinoma and sarcoma, respectively. While sarcomas are uncommon but most of the time highly aggressive, uterine carcinomas comprise the majority of cancer cases of the corpus uteri, but normally have a lower grade malignancy potential. More than 73% of the patients with the most frequent subtype of endometrial cancer present at FIGO stage I, resulting in a 5-y survival of 85 to 90%. However, more rare subtypes of endometrial carcinoma as well as uterine sarcoma display a more aggressive behavior and thus often present in more advanced stages with a 5-y survival of 5–20%, depending on the subtype. Surgery is the cornerstone of treatment. The value of adjuvant chemotherapy is controversial for carcinoma and proven not beneficial for high grade uterine sarcoma. In advanced or recurrent disease, treatment options are strongly limited.^{1,2} Therefore, there is an urgent need for exploration of new treatment options for uterine tumors.

One type of active immunotherapy is dendritic cell (DC) immunotherapy. DC have been discovered in 1973 by Ralph Steinman and since then, their role as sentinels of the

immune system that provide an essential link between innate and adaptive immunity has been elucidated. DC comprise only a minor (<1%) subset of the white blood cell population. Immature dendritic cells have the capacity to capture and process antigens, a process which leads to their maturation. Through interaction of the chemokine receptor CCR7 and its ligands CCL19/CCL21 DC migrate to the lymph nodes, where the processed antigens are presented to T cells, thus initiating an immune response.³ Since the isolation of circulating DCs is challenging, DC immunotherapy is most frequently based on DCs cultured in an ex vivo setting starting from CD34⁺ precursor cells or CD14⁺ monocytes. At the immature state DC are loaded with (defined) tumor antigens, offered as synthetic peptides or mRNA.

The application of active immunotherapy in the treatment of uterine cancer has not been explored for a long time and is still in its infancy, which is probably related to the poor insight into the interaction of the immune system with uterine cancers. However, as our knowledge increases, we believe that immunotherapeutic approaches hold promise for uterine cancer. Active immunotherapy relies on anticancer vaccines that are able to elicit an immune response against tumor-associated antigens (TAAs) in the human body. **Table 1** gives an overview of existing studies.

Loading of Tumor Antigens

Immunotherapeutic targets can consist of one or more defined TAA or a tumor-derived mixture of unknown TAA. Several TAA are validated for immunotherapy in uterine tumors such as for example several cancer testis antigens,⁸ MUC1,⁹ universal TAA such as hTERT or antigens targeting tumor cell survival such as survivin (Vanderstraeten et al., unpublished data). Until now the majority of antigen-focused immunotherapeutic studies in general have used a single antigen as a target. However, more and more arguments are arising for settings using a combination of multiple antigens or the use of total tumor cell approach. There are 3 major advantages. First of all, by using a combination of several antigens, the risk

of immune escape is reduced. Second, by using multiple antigens, the group of tumors and hence the patient group that can be targeted with one treatment approach is much larger. Lastly, by including anti-apoptosis factors as a target, the survival of the tumor itself can be attacked as well. Nevertheless, the use of a total tumor cell approach (tumor lysate, total tumoral mRNA,...) can theoretically induce auto-immunity, even though a certain amount of autoimmunity is necessary for efficient tumor eradication. To date, no reports of this type of side-effects have been published in the context of uterine cancer.

A crucial component in any DC vaccine is the choice of the DC loading strategy. Immunotherapy using DC loaded with peptide(s) is easily feasible but has several disadvantages, such as HLA restriction, the lack of a broad repertoire of MHC class II binding peptides and a short duration of antigen presentation. The in vitro work of several research groups has suggested that RNA transfection of TAA is an effective, if not superior, method to generate immunostimulatory DCs. By using mRNA encoding the whole TAA, RNA transfection is independent of the patient's HLA-type.^{10,11} In addition, further modification of the coding sequence by adding lysosomal targeting sequences (e.g., DC-LAMP) leads to the presentation of antigenic peptides in the context of both HLA class I and class II.¹²⁻¹⁴

Our research group has proven the safety and feasibility of DC immunotherapy in large series of high-grade glioblastoma (DCs loaded with lysate)¹⁵ and in case reports of uterine and ovarian tumors (*WT1* mRNA-loaded DCs).^{5,6,16}

Maturation Status of DC

DC vaccines should be able to induce high avidity CD8⁺ T cells that express high levels of granzyme and perforin and are able to enter the tumor microenvironment to exert their function. The quality of the induced CD8⁺ T cells is influenced by the expression of co-stimulatory molecules, such as CD80, CD86, CD137, and CD252 on DC, the cytokine secretion pattern of DC and the type of CD4⁺ helper T cell response.¹⁷⁻¹⁹ DC maturation is one

of the crucial factors.^{20,21} Currently, the “golden standard” used to mature DC consists of a cocktail of pro-inflammatory cytokines (IL-1 β , IL-6, TNF α) and prostaglandin E₂.²² However, a whole scale of cytokines, all mimicking the in vivo danger signal, is available to mature DC, such as IRX-2 and CD137.^{23,24} Recent mouse data demonstrated, however, that maturation of DC solely by pro-inflammatory cytokines yielded DC that failed to efficiently direct effector T cell differentiation. Interestingly, DC matured in the presence of Toll like receptor (TLR) ligands were able to induce full T cell effector function and induce more potent immune responses.^{25,26}

More recent in the field of DC maturation is the use of TriMixco-electroporation (caTLR4 [constitutively active Toll-like receptor 4], CD40L, and CD70 mRNA). The combination of CD40L and caTLR4 electroporation mimics CD40 ligation^{27,28} and TLR4 signaling²⁹ of the DC and generates phenotypically mature, cytokine/chemokine-secreting DC, as has been shown for CD40 and TLR4 ligation through addition of soluble CD40L and lipopolysaccharide (LPS).³⁰ On the other hand, the introduction of CD70 into the DC provides a co-stimulatory signal to CD27⁺ naive T cells by inhibiting activation-induced cell death (AICD) of T cells and by supporting T cell proliferation.³¹ DC co-electroporated with TriMix and tumor antigens have been shown to be potent stimulators of anti-tumor immunity in melanoma patients.³²⁻³⁵ Providing the DCs with a maturation signal through mRNA electroporation offers several advantages. There is no need to pre-incubate the DCs for up to 48 h with soluble maturation signals like pro-inflammatory cytokines or TLR ligands to achieve DC activation, which can render the cells “exhausted” and inferior for vaccination purposes.³⁶ As a result, TriMix DC, which can be injected into the patient within a short period after electroporation, will mature and secrete most of their immunostimulatory cytokines and chemokines in situ. Furthermore, it has been postulated that maturation of DCs

in situ resembles more closely the physiologic process involved in response to pathogen infection and may therefore lead to enhanced T-cell immunity.³⁷

Tumor Microenvironment

The sole application of DC immunotherapy to target uterine cancer is, however, not sufficient because of tumor-induced immunosuppression. More and more attention is being brought to the infiltration of immune suppressive cells in the tumor microenvironment as a response to classical treatments and immunotherapeutic strategies. These immune suppressive cells hamper the establishment of an effective anti-tumor immune response. To our knowledge, this has been poorly studied in uterine cancer and results are often contradictory.

The role of tumor-associated macrophages (TAMs) in endometrial carcinoma is complex. Located in close contact with cancer cells, they have a beneficial effect on relapse-free survival, but TAMs located in necrotic tumor tissue are linked with tumor recurrence.³⁸ Moreover, TAMs secrete matrix metalloproteinases and induce angiogenesis which promote tumor progression.³⁹ Furthermore, TAMs are associated with myometrial invasion and lymph node metastasis and thus have a clear tumor-promoting role.^{40,41}

Endometrial tumors are frequently infiltrated by T lymphocytes as well. CD8⁺ lymphocytes present in primary untreated stage IA-IIIa endometrial cancer, show upregulation of inhibitory natural killer receptors, an effect that possibly is mediated through TGF- β . However, this upregulation prevents the cytotoxic function of these lymphocytes at the HLA recognition level, thus causing an important tumor evasion strategy, since non-classical HLA molecules are upregulated in cancer.⁴² A recent publication showed the association of uterine sarcoma with systemic inflammation, as measured by an enhanced neutrophil to lymphocyte ratio in these patients.⁴³ Tumor-infiltrating lymphocytes (TIL) can be detected in uterine sarcoma, but these cells can sometimes harbor chromosomal aberrations which might negatively impact their functionality.⁴⁴

In many tumors, regulatory T cells (Treg) suppress endogenous and induced antitumor immune responses. However, in endometrial carcinoma, the presence of Treg is debatable. Fattorossiet al.⁴⁵ showed their presence in the tumor-draining lymph nodes of 26 patients, whereas Giatromanolakiet al.⁴⁶ demonstrated less Treg in 79 stage I endometrial carcinoma patients compared with normal endometrium. Recent work of both Yamagamiet al.⁴⁷ and Zhang et al.⁴⁸ demonstrated an increase of Treg in the tumor tissue and peripheral blood respectively in endometrial carcinoma, increasing in more advanced stages. In contrast, the group of Sawanet al.⁴⁹ attributed this increase in Tregs to an increase in age and to the postmenopausal condition of most patients.

More recently, myeloid derived suppressor cells (MDSC) have been shown to play an important role in cancer immunosuppression and have been implicated in several tumor types. We showed for the first time that both endometrial carcinoma and uterine sarcoma tumors contain MDSC infiltrates that express arginase-1.⁵⁰

Proof of Concept

In order to improve our previous DC vaccination regimen, several modifications were incorporated:

- 1) Inclusion of both WT1 and survivin as targets to broaden the immune response
- 2) Inclusion of the DC-LAMP targeting signal in the WT1 and survivin mRNA for induction of both CD4⁺ and CD8⁺ T cells
- 3) Co-electroporation of WT1 and survivin with TriMix to induce DC maturation.

The feasibility of this approach was tested on leukapheresis products of 3 healthy donors. Monocytes were isolated by elutriation and cultured in CellGenix culture bags in the presence of GM-CSF and IL-4 (Cellgenix) until day 7. Immature DC were harvested, electroporated with mRNA encoding WT1-DC.LAMP and TriMix or survivin-DC.LAMP and

TriMix, mixed and cultured for 3.5 to 4 h in the presence of rapamycin (MiltenyiBiotec Leiden), IL-4, and GM-CSF before cryopreservation.

Upon thawing, DC showed a viability of $57.3 \pm 5.5\%$ and a purity of $64.6 \pm 7.8\%$. DC phenotype was analyzed after thawing (4h after electroporation) and after an additional 44h culture period (48h after electroporation). Immediately after thawing, DC still presented with a rather immature phenotype, but became fully mature at 48h after electroporation, with upregulation of CD54, CD80, CD83, CD86, and CCR7 (**Table 2**). WT1 expression by DC was at the highest level 1.5 h after thawing and steadily decreased thereafter, while survivin only slightly decreased at 48h after thawing (**Fig. 1**). DC secreted low levels of IL-12p70, intermediate levels of IL-1 β and IL-18 and high levels of IL-6, IL-8, MCP-1, and MIP-1 α . Upon stimulation of autologous T cells, we observed an increase of survivin-tetramer (survivin 95–104 peptide) positive T cells from almost undetectable (0.08% of CD8⁺ T cells) to 0.56% of CD8⁺ T cells after 1 stimulation (7-fold) and to 0.81% of CD8⁺ T cells after 2 stimulations with DC (10-fold). No induction of WT1-specific T cells was noted (**Fig. 2**).

Conclusions

DC immunotherapy is in its infancy in uterine cancer. Nevertheless, the recent data concerning the interaction of uterine tumors with the immune system indicate that immunotherapy is promising as a novel therapeutic approach for this type of cancer. Furthermore, immunotherapeutic approaches are associated with less toxicity and fewer side effects, which is important for an elderly patient population. We have previously shown the feasibility of DC vaccination in uterine cancer and we anticipate that our novel approach will further improve this modality. Moreover, in our opinion, the combination of DC vaccines with inhibitors of immunosuppression holds great promise for uterine cancer patients.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The Authors would like to thank Kim Morias of Immunotherapy Platform Leuven (www.itpl.be) for the manufacturing of the dendritic cells.

A.C. is supported by the Fund for Scientific Research (FWO-Vlaanderen). A.V. and S.T. are supported by the Verelst Uterine Cancer Fund Leuven. S.W.V.G. and F.A. are Senior Clinical Investigators of the Fund for Scientific Research–Flanders (FWO-Vlaanderen)

This project is supported by the Olivia Hendrickx Research Fund (www.olivia.be), by the Anticancer Fund (www.reliablecancertherapies.com) and by the Research Fund of the University Hospital Leuven.

A.C. is a postdoctoral research fellow supported by the FWO-V; F.A. and S.W.VG. are senior clinical investigators supported by the FWO-V.

References

- <jrn>1. AmantF, MoermanP, NevenP, TimmermanD, Van LimbergenE, VergoteI. Endometrial cancer.Lancet2005; 366:491-505; PMID:16084259; [http://dx.doi.org/10.1016/S0140-6736\(05\)67063-8](http://dx.doi.org/10.1016/S0140-6736(05)67063-8)</jrn>
- <jrn>2. AmantF, CoosemansA, Debiec-RychterM, TimmermanD, VergoteI. Clinical management of uterine sarcomas.Lancet Oncol2009; 10:1188-98; PMID:19959075; [http://dx.doi.org/10.1016/S1470-2045\(09\)70226-8](http://dx.doi.org/10.1016/S1470-2045(09)70226-8)</jrn>
- <jrn>3. BanchereauJ, BriereF, CauxC, DavoustJ, LebecqueS, LiuYJ, PulendranB, PaluckaK. Immunobiology of dendritic cells.Annu Rev Immunol2000; 18:767-811; PMID:10837075; <http://dx.doi.org/10.1146/annurev.immunol.18.1.767></jrn>
- <jrn>4. SantinAD, BelloneS, RavaggiA, RomanJJ, PecorelliS, ParhamGP, CannonMJ. Induction of tumour-specific CD8(+) cytotoxic T lymphocytes by tumour lysate-pulsed autologous dendritic cells in patients with uterine serous papillary cancer.Br J Cancer2002; 86:151-7; PMID:11857027; <http://dx.doi.org/10.1038/sj.bjc.6600026></jrn>
- <jrn>5. HernandoJJ, ParkTW, KüblerK, OffergeldR, SchlebuschH, BauknechtT. Vaccination with autologous tumour antigen-pulsed dendritic cells in advanced gynaecological malignancies: clinical and immunological evaluation of a phase I trial.Cancer ImmunolImmunother2002; 51:45-52; PMID:11845259; <http://dx.doi.org/10.1007/s00262-001-0255-1></jrn>

- <jrn>6. CoosemansA, WölflM, BernemanZN, Van TendelooV, VergoteI, AmantF, Van GoolSW. Immunological response after therapeutic vaccination with WT1 mRNA-loaded dendritic cells in end-stage endometrial carcinoma. *Anticancer Res*2010; 30:3709-14; PMID:20944158</jrn>
- <jrn>7. CoosemansA, VanderstraetenA, TuyartsS, VerschuerT, MoermanP, BernemanZN, VergoteI, AmantF, VAN GoolSW. Wilms' Tumor Gene 1 (WT1)--loaded dendritic cell immunotherapy in patients with uterine tumors: a phase I/II clinical trial. *Anticancer Res*2013; 33:5495-500; PMID:24324087</jrn>
- <jrn>8. BrooksN, PouniotisDS. Immunomodulation in endometrial cancer. *Int J Gynecol Cancer*2009; 19:734-40; PMID:19509580; <http://dx.doi.org/10.1111/IGC.0b013e3181a12f7f></jrn>
- <jrn>9. SivridisE, GiatromanolakiA, KoukourakisMI, GeorgiouL, AnastasiadisP. Patterns of episialin/MUC1 expression in endometrial carcinomas and prognostic relevance. *Histopathology*2002; 40:92-100; PMID:11903603; <http://dx.doi.org/10.1046/j.1365-2559.2002.01316.x></jrn>
- <jrn>10. GilboaE. DC-based cancer vaccines. *J Clin Invest*2007; 117:1195-203; PMID:17476349; <http://dx.doi.org/10.1172/JCI31205></jrn>
- <jrn>11. MichielsA, TuyartsS, BonehillA, CorthalsJ, BreckpotK, HeirmanC, Van MeirvenneS, DullaersM, AllardS, BrasseurF, et al. Electroporation of immature and mature dendritic cells: implications for dendritic cell-based vaccines. *Gene Ther*2005; 12:772-82; PMID:15750615; <http://dx.doi.org/10.1038/sj.gt.3302471></jrn>
- <jrn>12. BonehillA, HeirmanC, ThielemansK. Genetic approaches for the induction of a CD4+ T cell response in cancer immunotherapy. *J Gene Med*2005; 7:686-95; PMID:15693037; <http://dx.doi.org/10.1002/jgm.713></jrn>
- <jrn>13. BonehillA, HeirmanC, TuyartsS, MichielsA, BreckpotK, BrasseurF, ZhangY, Van Der BruggenP, ThielemansK. Messenger RNA-electroporated dendritic cells presenting MAGE-A3 simultaneously in HLA class I and class II molecules. *J Immunol*2004; 172:6649-57; PMID:15153480</jrn>
- <jrn>14. SuZ, DannullJ, YangBK, DahmP, ColemanD, YanceyD, SichiS, NiedzwieckiD, BoczkowskiD, GilboaE, et al. Telomerase mRNA-transfected dendritic cells stimulate antigen-specific CD8+ and CD4+ T cell responses in patients with metastatic prostate cancer. *J Immunol*2005; 174:3798-807; PMID:15749921</jrn>
- <jrn>15. ArdonH, Van GoolSW, VerschuerT, MaesW, FieuwsS, SciotR, WilmsG, DemareelP, GoffinJ, VanCalenberghF, et al. Integration of autologous dendritic cell-based immunotherapy in the standard of care treatment for patients with newly diagnosed glioblastoma: results of the HGG-2006 phase I/II trial. *Cancer Immunol Immunother*2012; 61:2033-44; PMID:22527250; <http://dx.doi.org/10.1007/s00262-012-1261-1></jrn>
- <jrn>16. CoosemansA, VanderstraetenA, TuyartsS, VerschuerT, MoermanP, BernemanZ, VergoteI, AmantF, Van GoolSW. Immunological response after WT1 mRNA-loaded dendritic cell immunotherapy in ovarian carcinoma and carcinosarcoma. *Anticancer Res*2013; 33:3855-9; PMID:24023319</jrn>
- <jrn>17. PaluckaK, BanchereauJ. Cancer immunotherapy via dendritic cells. *Nat Rev Cancer*2012; 12:265-77; PMID:22437871; <http://dx.doi.org/10.1038/nrc3258></jrn>
- <jrn>18. PaluckaK, UenoH, BanchereauJ. Recent developments in cancer vaccines. *J Immunol*2011; 186:1325-31; PMID:21248270; <http://dx.doi.org/10.4049/jimmunol.0902539></jrn>
- <jrn>19. Van BrusselJ, BernemanZN, CoolsN. Optimizing dendritic cell-based immunotherapy: tackling the complexity of different arms of the immune system. *Mediators Inflamm*2012; 2012:690643; PMID:22851815; <http://dx.doi.org/10.1155/2012/690643></jrn>

- <jrn>20. De VriesIJ, KrooshoopDJ, ScharenborgNM, LesterhuisWJ, DiepstraJH, Van MuijenGN, StrijkSP, RuersTJ, BoermanOC, OyenWJ, et al. Effective migration of antigen-pulsed dendritic cells to lymph nodes in melanoma patients is determined by their maturation state. *Cancer Res*2003; 63:12-7; PMID:12517769</jrn>
- <jrn>21. de VriesIJ, LesterhuisWJ, BarentszJO, VerdijkP, van KriekenJH, BoermanOC, OyenWJ, BonenkampJJ, BoezemanJB, AdemaGJ, et al. Magnetic resonance tracking of dendritic cells in melanoma patients for monitoring of cellular therapy. *Nat Biotechnol*2005; 23:1407-13; PMID:16258544; <http://dx.doi.org/10.1038/nbt1154></jrn>
- <jrn>22. JonuleitH, KühnU, MüllerG, SteinbrinkK, ParagnikL, SchmittE, KnopJ, EnkAH. Pro-inflammatory cytokines and prostaglandins induce maturation of potent immunostimulatory dendritic cells under fetal calf serum-free conditions. *Eur J Immunol*1997; 27:3135-42; PMID:9464798; <http://dx.doi.org/10.1002/eji.1830271209></jrn>
- <jrn>23. SchillingB, HarasymczukM, SchulerP, EganJ, FerroneS, WhitesideTL. IRX-2, a novel immunotherapeutic, enhances functions of human dendritic cells. *PLoS One*2013; 8:e47234; PMID:23408925; <http://dx.doi.org/10.1371/journal.pone.0047234></jrn>
- <jrn>24. HarfuddinZ, KwajahS, Chong NyiSimA, MacaryPA, SchwarzH. CD137L-stimulated dendritic cells are more potent than conventional dendritic cells at eliciting cytotoxic T-cell responses. *Oncoimmunology*2013; 2:e26859; PMID:24482752; <http://dx.doi.org/10.4161/onci.26859></jrn>
- <jrn>25. SpörriR, Reis e SousaC. Inflammatory mediators are insufficient for full dendritic cell activation and promote expansion of CD4+ T cell populations lacking helper function. *Nat Immunol*2005; 6:163-70; PMID:15654341; <http://dx.doi.org/10.1038/ni1162></jrn>
- <jrn>26. MailliardRB, Wankowicz-KalinskaA, CaiQ, WesaA, HilkenCM, KapsenbergML, KirkwoodJM, StorkusWJ, KalinskiP. alpha-type-1 polarized dendritic cells: a novel immunization tool with optimized CTL-inducing activity. *Cancer Res*2004; 64:5934-7; PMID:15342370; <http://dx.doi.org/10.1158/0008-5472.CAN-04-1261></jrn>
- <jrn>27. KikuchiT, MooreMA, CrystalRG. Dendritic cells modified to express CD40 ligand elicit therapeutic immunity against preexisting murine tumors. *Blood*2000; 96:91-9; PMID:10891436</jrn>
- <jrn>28. KikuchiT, WorgallS, SinghR, MooreMA, CrystalRG. Dendritic cells genetically modified to express CD40 ligand and pulsed with antigen can initiate antigen-specific humoral immunity independent of CD4+ T cells. *Nat Med*2000; 6:1154-9; PMID:11017148; <http://dx.doi.org/10.1038/80498></jrn>
- <jrn>29. CiscoRM, Abdel-WahabZ, DannullJ, NairS, TylerDS, GilboaE, ViewegJ, DaakaY, PruittSK. Induction of human dendritic cell maturation using transfection with RNA encoding a dominant positive toll-like receptor 4. *J Immunol*2004; 172:7162-8; PMID:15153540</jrn>
- <jrn>30. LapointeR, TosoJF, ButtsC, YoungHA, HwuP. Human dendritic cells require multiple activation signals for the efficient generation of tumor antigen-specific T lymphocytes. *Eur J Immunol*2000; 30:3291-8; PMID:11093145; [http://dx.doi.org/10.1002/1521-4141\(200011\)30:11<3291::AID-IMMU3291>3.0.CO;2-2](http://dx.doi.org/10.1002/1521-4141(200011)30:11<3291::AID-IMMU3291>3.0.CO;2-2)</jrn>
- <jrn>31. BorstJ, HendriksJ, XiaoY. CD27 and CD70 in T cell and B cell activation. *Curr Opin Immunol*2005; 17:275-81; PMID:15886117; <http://dx.doi.org/10.1016/j.coi.2005.04.004></jrn>
- <jrn>32. WilgenhofS, Van NuffelAM, CorthalsJ, HeirmanC, TuyaertsS, Benteynd, De ConinckA, Van RietI, VerfaillieG, VandeloofJ, et al. Therapeutic vaccination with an autologous mRNA electroporated dendritic cell

- vaccine in patients with advanced melanoma. *J Immunother* 2011; 34:448-56; PMID:21577140; <http://dx.doi.org/10.1097/CJI.0b013e31821dcb31></jrn>
- <jrn>33. Bonehill A, Tuyaerts S, Van Nuffel AM, Heirman C, Bos TJ, Fostier K, Neyns B, Thielemans K. Enhancing the T-cell stimulatory capacity of human dendritic cells by co-electroporation with CD40L, CD70 and constitutively active TLR4 encoding mRNA. *Mol Ther* 2008; 16:1170-80; PMID:18431362; <http://dx.doi.org/10.1038/mt.2008.77></jrn>
- <jrn>34. Bonehill A, Van Nuffel AM, Corthals J, Tuyaerts S, Heirman C, François V, Colau D, van der Bruggen P, Neyns B, Thielemans K. Single-step antigen loading and activation of dendritic cells by mRNA electroporation for the purpose of therapeutic vaccination in melanoma patients. *Clin Cancer Res* 2009; 15:3366-75; PMID:19417017; <http://dx.doi.org/10.1158/1078-0432.CCR-08-2982></jrn>
- <jrn>35. Van Nuffel AM, Tuyaerts S, Benteyn D, Wilgenhof S, Corthals J, Heirman C, Neyns B, Thielemans K, Bonehill A. Epitope and HLA-type independent monitoring of antigen-specific T-cells after treatment with dendritic cells presenting full-length tumor antigens. *J Immunol Methods* 2012; 377:23-36; PMID:22269772; <http://dx.doi.org/10.1016/j.jim.2011.12.010></jrn>
- <jrn>36. Langenkamp A, Messi M, Lanzavecchia A, Sallusto F. Kinetics of dendritic cell activation: impact on priming of TH1, TH2 and nonpolarized T cells. *Nat Immunol* 2000; 1:311-6; PMID:11017102; <http://dx.doi.org/10.1038/79758></jrn>
- <jrn>37. Nair S, McLaughlin C, Weizer A, Su Z, Boczkowski D, Dannull J, Vieweg J, Gilboa E. Injection of immature dendritic cells into adjuvant-treated skin obviates the need for ex vivo maturation. *J Immunol* 2003; 171:6275-82; PMID:14634145</jrn>
- <jrn>38. Ohno S, Ohno Y, Suzuki N, Kamei T, Koike K, Inagawa H, Kohchi C, Soma G, Inoue M. Correlation of histological localization of tumor-associated macrophages with clinicopathological features in endometrial cancer. *Anticancer Res* 2004; 24(5C):3335-42; PMID:15515429</jrn>
- <jrn>39. Iurlaro M, Loverro G, Vacca A, Cormio G, Ribatti D, Minischetti M, Ria R, Bruno M, Selvaggi L. Angiogenesis extent and expression of matrix metalloproteinase-2 and -9 correlate with upgrading and myometrial invasion in endometrial carcinoma. *Eur J Clin Invest* 1999; 29:793-801; PMID:10469168; <http://dx.doi.org/10.1046/j.1365-2362.1999.00532.x></jrn>
- <jrn>40. Soeda S, Nakamura N, Ozeki T, Nishiyama H, Hojo H, Yamada H, Abe M, Sato A. Tumor-associated macrophages correlate with vascular space invasion and myometrial invasion in endometrial carcinoma. *Gynecol Oncol* 2008; 109:122-8; PMID:18289648; <http://dx.doi.org/10.1016/j.ygyno.2007.12.033></jrn>
- <jrn>41. Espinosa I, José Carnicer M, Catusus L, Canet B, D'angelo E, Zannoni GF, Prat J. Myometrial invasion and lymph node metastasis in endometrioid carcinomas: tumor-associated macrophages, microvessel density, and HIF1A have a crucial role. *Am J Surg Pathol* 2010; 34:1708-14; PMID:20962622</jrn>
- <jrn>42. Chang WC, Huang SC, Torng PL, Chang DY, Hsu WC, Chiou SH, Chow SN, Sheu BC. Expression of inhibitory natural killer receptors on tumor-infiltrating CD8+ T lymphocyte lineage in human endometrial carcinoma. *Int J Gynecol Cancer* 2005; 15:1073-80; PMID:16343184; <http://dx.doi.org/10.1111/j.1525-1438.2005.00264.x></jrn>

- <jrn>43. Kim HS, Han KH, Chung HH, Kim JW, Park NH, Song YS, Kang SB. Neutrophil to lymphocyte ratio for preoperative diagnosis of uterine sarcomas: a case-matched comparison. *Eur J Surg Oncol* 2010; 36:691-8; PMID:20570475; <http://dx.doi.org/10.1016/j.ejso.2010.05.004></jrn>
- <jrn>44. Engel H, Friedrich I, Kleespies C, Kurbacher CM, Schöndorf T, Grecu O, Kolhagen H, Mallmann P. Detection of chromosomal aberrations in tumor cells and tumor infiltrating lymphocytes by molecular cytogenetics in patients with gynecological cancer. *Cancer Genet Cytogenet* 1998; 106:159-65; PMID:9797783; [http://dx.doi.org/10.1016/S0165-4608\(98\)00070-3](http://dx.doi.org/10.1016/S0165-4608(98)00070-3)</jrn>
- <jrn>45. Fattorossi A, Battaglia A, Ferrandina G, Buzzonetti A, Legge F, Salutari V, Scambia G. Lymphocyte composition of tumor draining lymph nodes from cervical and endometrial cancer patients. *Gynecol Oncol* 2004; 92:106-15; PMID:14751146; <http://dx.doi.org/10.1016/j.ygyno.2003.09.020></jrn>
- <jrn>46. Giatromanolaki A, Bates GJ, Koukourakis MI, Sivridis E, Gatter KC, Harris AL, Banham AH. The presence of tumor-infiltrating FOXP3+ lymphocytes correlates with intratumoral angiogenesis in endometrial cancer. *Gynecol Oncol* 2008; 110:216-21; PMID:18533240; <http://dx.doi.org/10.1016/j.ygyno.2008.04.021></jrn>
- <jrn>47. Yamagami W, Susumu N, Tanaka H, Hirasawa A, Banno K, Suzuki N, Tsuda H, Tsukazaki K, Aoki D. Immunofluorescence-detected infiltration of CD4+FOXP3+ regulatory T cells is relevant to the prognosis of patients with endometrial cancer. *Int J Gynecol Cancer* 2011; 21:1628-34; PMID:21897268; <http://dx.doi.org/10.1097/IGC.0b013e31822c271f></jrn>
- <jrn>48. Zhang W, Hou F, Zhang Y, Tian Y, Jiao J, Ma D, Kong B, Cui B. Changes of Th17/Tc17 and Th17/Treg cells in endometrial carcinoma. *Gynecol Oncol* 2014; 132:599-605; PMID:24388919; <http://dx.doi.org/10.1016/j.ygyno.2013.12.036></jrn>
- <jrn>49. Sawan S, Burt DJ, Stern PL, Holland C, Elkord E. Circulating regulatory T cells in endometrial cancer: a role for age and menopausal status. *Immunol Invest* 2011; 40:62-75; PMID:20809698; <http://dx.doi.org/10.3109/08820139.2010.513022></jrn>

50. Vanderstraeten A, Luyten C, Verbist G, Tuyaerts S, Amant F. Mapping the immunosuppressive environment in uterine tumors: implications for immunotherapy. *Cancer Immunol Immunother* 2014

Figure 1. WT1 or survivin antigen expression by mRNA electroporated DC. DC electroporated with WT1.DC-LAMP + TriMix or survivin.DC-LAMP + TriMix were analyzed at 4h or 48h after electroporation for expression of WT1 or survivin by immunocytochemistry. The percentage of stained cells (left panel) and the intensity of staining (right panel) was documented.




Figure 2. Tetramer analysis of stimulated T cells. DC electroporated with WT1.DC-LAMP + TriMix or survivin.DC-LAMP + TriMix were used at 4h post electroporation to stimulate autologous T cells. Unstimulated T cells and stimulated T cells were stained for CD3, CD8, and the respective tetramer. Cells were gated on FSC/SSC characteristics and CD3⁺ CD8⁺ cells. On the CD3⁺ CD8⁺ T cells, the percentage of WT1- or survivin-tetramer positive cells was determined.




Table 1. Overview on DC immunotherapy in uterine cancer

Group	Tumor	Number of Patients	Type of Active Immunotherapy
Santin A et al. ⁴ 2002	Serous endometrial carcinoma	3	Injection of DCs loaded with whole tumor lysate
Hernando JJ et al. ⁵ 2002	Uterine sarcoma	2	Injection of DCs loaded with whole tumoral lysate and KLH
Coosemans A et al. ⁶ 2010	Serous endometrial carcinoma	1	WT1-mRNA loaded DC immunotherapy
Coosemans A et al. ⁷ 2013	Leiomyosarcoma and serous endometrial carcinoma	6	WT1-mRNA loaded DC immunotherapy

KLH, Keyhole Limpet Hemocyanin; WT1, Wilms' tumor gene 1; DC: dendritic cell.

Table 2. Characteristics of DC from 3 test leukaphereses (n = 3)

Test	4 hours	48 hours
Sterility		
bacteria	negative	negative

mycoplasma	negative	negative
fungi	negative	negative
endotoxins	negative	negative
mDC-identity*		
CD11c	90.9% (12.7)	95.4% (2.2)
CD14	11.8% (8.3)	5.3% (5.3)
CD54	89.1% (11.8)	98.0% (3.7)
CD80	38.1% (15.5)	98.3% (1.0)
CD83	30.4% (8.0)	65.4% (9.0)
CD86	76.7% (18.4)	98.3% (0.6)
CCR7	25.8% (11.4)	96.5% (2.0)
HLA-ABC	99.0% (0.6)	99.7% (0.2)
HLA-DR	98.1% (2.2)	97.6% (2.1)
mRNA		
electroporation*		
CD70	45.5% (11.8)	57.9% (22.8)
Viability upon		
thawing*		
	57.3% (5.5)	57.3% (5.5)
Purity upon		
thawing*		
	64.6% (7.8)	64.6% (7.8)

*Results expressed as mean percentage (standard deviation).